



540 DIVISION STREET = CAMPBELL = CALIFORNIA 95008-6906 = USA 408-866-6363 = 800-726-3213 = FAX 408-866-6364 = EMAIL info@listlabs.com WEBSITE www.listlabs.com

# CERTIFICATE OF ANALYSIS Enterotoxin type B from Staphylococcus aureus Lot #1223B1

# Contents:

Each vial, when reconstituted with 1.0 ml of sterile distilled water, contains 0.5 mg of enterotoxin type B in 0.005 M potassium phosphate, pH 6.8.

# Method of Preparation:

This product is prepared from *S. aureus* strain S6 by modifications of the ion exchange chromatography method of Schantz, *et al.*<sup>1</sup>

### Concentration:

Protein concentration is determined using an extinction coefficient of  $E_{1 \text{ cm}}^{1 \text{ mg/ml}} = 1.44$  measured at 277 nm.<sup>2</sup>

### Electrophoretic Purity:

This preparation migrates as a single major band of molecular weight approximately 28,000 daltons when run on 12% SDS-polyacrylamide gels prepared according to the method of Wyckoff,<sup>3</sup> a modification of the Laemmli<sup>4</sup> gel system.

# Packaging and Storage:

This preparation is filter sterilized using a 0.2  $\mu$  filter and packaged under aseptic conditions. It is supplied as a lyophilized powder, sealed under vacuum. Store at 4°C prior to reconstitution. Following reconstitution, maintain sterility and store frozen in aliquots.

### Toxicity:

The emetic dose 50% in animals is approximately 1  $\mu$ g/kg intragastrically, and 0.1 to 0.5  $\mu$ g/kg intravenously.<sup>5</sup> Humans are more sensitive, and it is estimated that 2-3 ng/kg causes illness. Enterotoxin type B may be lethal at high doses. This product is a select agent.

(continued)

# **Handling:**

Good laboratory technique should be employed in the safe handling of this product. This involves observing the following practices:

- 1. This product is to be used by skilled personnel under the direction of a principal investigator in a biosafety level 2 containment area only. Biosafety level 3 precautions may be indicated for activities with a high potential for aerosol or droplet production.
- 2. Wear appropriate attire, i.e., lab coat, eye protection and latex gloves.
- 3. <u>Never</u> remove the stopper prior to reconstitution, and <u>never</u> work with the product in the powdered form. Always reconstitute it first.
- 4. Do not mouth pipette, inhale, ingest or allow to come into contact with open wounds. Wash thoroughly any area of the body which comes into contact with the product.
- 5. Avoid accidental autoinoculation by exercising extreme care when handling in conjunction with any injection device.
- 6. This product is intended for research purposes only. It is not intended for use in humans or as a diagnostic agent. List Biological Laboratories, Inc. is not liable for any damages resulting from the misuse or handling of this product.

# FOR RESEARCH PURPOSES ONLY. NOT FOR HUMAN USE.

### References:

- 1. Schantz, E.J., Roessler, W.G., Wagman, J., Spero, L., Dunnery, D.A., and Bergdoll, M.S. (1965) *Biochemistry* **4 (6)**, 1011-1016.
- 2. landolo, J.J. and Tweten, R.K. (1988) Meth. Enzymol. 165, 43-52.
- 3. Wyckoff, M., Rodbard, D. and Chrambach, A. (1977) *Anal. Biochem.* **78**, 459-482.
- 4. Laemmli, U.K. (1970) Nature 227, 680-685.
- 5. Bergdoll, M.S. (1988) *Meth. Enzymol.* **165**, 324-333.

| Approved: | KA | Date:_ | 5/26 | 106 | Approved: | M | Date: 5/30/04 |
|-----------|----|--------|------|-----|-----------|---|---------------|
|           |    |        | 1 1  |     |           |   |               |