



540 Division Street, Campbell, California 95008-6906  
Tel. (408) 866-6363 Fax (408) 866-6364  
[www.listlabs.com](http://www.listlabs.com)

Product #172B  
Lot #1722B15A1  
Release Date: July 2017

**CERTIFICATE OF ANALYSIS  
ANTHRAX LETHAL FACTOR (LF)  
RECOMBINANT  
from *Bacillus anthracis*  
Lot #1722B15A1**

**Contents**

Reconstitute with 0.5 mL of water. Each vial contains 1 mg of lethal factor from *Bacillus anthracis* in 5 mM HEPES, 50 mM NaCl, pH 7.5. **Handle the product gently. Do not vortex.**

**Packaging/Storage**

This product is packaged aseptically, lyophilized, and sealed under vacuum. Store at -20°C prior to reconstitution.

**Concentration**

Protein concentration was determined by absorbance at 280 nm using an extinction coefficient of 0.88 for a 1 mg/mL solution. This value is calculated by ProtParam<sup>1</sup> using an algorithm based on the Edelhoch<sup>2</sup> method with modifications described in Pace et al.<sup>3</sup>

Protein concentration was also confirmed by a modification of Bradford<sup>4</sup>, using NIST traceable bovine serum albumin as the standard.

**Purity**

When examined on 4 – 12% SDS-PAGE gels, this protein migrates as a single major band with an apparent molecular weight of approximately 90,000 daltons. Densitometric analysis estimates the purity of the product as  $\geq 90\%$ .

The endotoxin content, determined using a kinetic chromogenic LAL assay, is  $< 2$  EU/mg.

**Activity**

LF is assessed for specific activity in units/mg protein, using MAPKKide<sup>®</sup>, Product #530, which is a synthetic FRET peptide containing a single cleavage site for LF. A standard curve generated from MAPKKide<sup>®</sup> Unquenched Calibration Peptide for #530, Product #539, is used to convert relative fluorescence units (RFU) to  $\mu$ moles of cleaved substrate. One unit of lethal factor will catalyze the release of 1.0  $\mu$ mole of cleaved MAPKKide<sup>®</sup> per minute at 37°C in 20 mM HEPES, pH 8.2. The specific activity of this lot of lethal factor is 0.3 units/mg.

LF is assessed for cytotoxicity in the presence of 1  $\mu$ g/mL PA using J774A.1 cells. When J774A.1 cells are treated with LF alone, no toxicity was seen at 1  $\mu$ g/mL (11 nM). The effective concentration 50% (EC<sub>50</sub>) of this LF lot meets specifications.



540 Division Street, Campbell, California 95008-6906  
Tel. (408) 866-6363 Fax (408) 866-6364  
[www.listlabs.com](http://www.listlabs.com)

Product #172B  
Lot #1722B15A1  
Release Date: July 2017

### Handling

Good laboratory technique should be employed in the safe handling of this product. Wear appropriate laboratory attire including lab coat, gloves, and safety glasses. Nitrile gloves are recommended when handling lyophilized material.

This product is intended for research purposes by qualified personnel. It is not intended for use in humans or as a diagnostic agent. List Biological Laboratories, Inc. is not liable for any damages resulting from the misuse or handling of this product.

**FOR RESEARCH PURPOSES ONLY. NOT FOR HUMAN USE.**

### References

1. [www.expasy.ch/tools/protparam-doc.html](http://www.expasy.ch/tools/protparam-doc.html)
2. Edelhoch, H. (1967) *Biochemistry*, **6**, 1948-1954.
3. Pace, C.N., Vajdos, F., Fee, L., Grimsley, G. and Gray, T. (1995) *Protein Sci.* **4**, 2411-2423.
4. Bradford, M.M. (1976) *Anal. Biochem.* **72**, 248-254.

Research:  Date: 09/03/2021