

540 Division Street, Campbell, California 95008-6906 Tel. (408) 866-6363 Fax (408) 866-6364 www.listlabs.com

Product #190A Lot #19050A1

Release Date: Sept 2019

CERTIFICATE OF ANALYSIS TETANUS TOXIN from Clostridium tetani Lot #19050A1



Contents

Each vial contains 25 µg of tetanus toxin. When reconstituted to 100 µL with water, the buffer is approximately 20 mM HEPES, pH 7.4 with 1.25% lactose. Handle the product gently; do not vortex.

Concentration

The protein concentration was determined by absorbance at 280 nm using an extinction coefficient of 1.26 for a 1 mg/ml solution. This value is calculated by ProtParam¹ using an algorithm based on the Edelhoch² method with modifications described in Pace et al³.

NOTE: The concentration for previous lots of Product #190A was determined by a modification of the method of Bradford⁴ using ovalbumin as the standard.

Purity

When examined on 4 – 12% SDS-PAGE, this protein migrates as a single major band with an apparent molecular weight of approximately 150,000 daltons. In the presence of a reducing agent, the preparation migrates as two bands with apparent molecular weights of 100,000 and 50,000 daltons. Densitometric analysis estimates the purity as >80%.

The endotoxin content, determined using a kinetic chromogenic LAL assay, is 2 EU/mg.

Activity

Binding activity to G_{T1b} ganglioside in a hemagglutination assay is also assessed.⁵ Hemagglutination is evident at 6.25 µg/ml Tetanus Toxin.

The tetanus toxin has been tested for enzymatic activity in a FRET based assay. Digestion of the FRET substrate was detected within 3 hours when using 200 nM tetanus toxin with 20 µM VAMPtide (Product #540).

Toxicity

Tetanus toxin is one of the most deadly toxins known to man. Even small amounts of tetanus toxin can pose a serious threat to an unvaccinated user. Consult the SDS for further information.

Packaging/Storage

This preparation is provided as a lyophilized powder that has been stoppered under vacuum. Prior to reconstitution, it should be stored at 2 - 8°C.

Handling

Good laboratory technique should be employed in the safe handling of this product. Wear appropriate laboratory attire including a lab coat, gloves and safety glasses. Nitrile gloves are recommended for use when handling lyophilized material.

This product is intended for research purposes by qualified personnel. It is not intended for use in humans or as a diagnostic agent. List Biological Laboratories, Inc., is not liable for any damages resulting from the misuse or handling of this product.

FOR RESEARCH PURPOSES ONLY. NOT FOR HUMAN USE

References

- 1. www.expasy.ch/tools/protparam-doc.html
- 2. Edelhoch, H. (1967) Biochemistry, 6, 1948-1954.
- 3. Pace, C.N., Vajdos, F., Fee, L., Grimsley, G. and Gray, T. (1995) Protein Sci. 4, 2411-2423.
- 4. Bradford, M.M. (1976) Anal. Biochem. 72, 248 254.
- 5. Tayot, J.-L., Holmgren, J., Svennerholm L., Lindblad, M. and Tardy, M. (1981) Eur. J. Biochem. 113, 249 - 258.

Recommendations

List Labs recommends that any person who may potentially become exposed to this Toxin should be currently vaccinated with tetanus toxoid.

The Toxin should be controlled and maintained in full compliance with biosafety level 2 or 3 standards, whichever is required based on customer's utilization. Such standards are described in "Biosafety in Microbiology and Biomedical Laboratories" HHS Publication No. (NIH) 88-8395, 2nd edition, May 1988 and as the same may be amended or superseded.

Personnel who come into contact with the Toxin shall be fully trained and qualified in its handling and use.

QA/QC: KPP Date: 28 JAN 30 21

