



**LIST BIOLOGICAL  
LABORATORIES, INC.**

540 DIVISION STREET CAMPBELL CALIFORNIA 95008-6906 USA  
408-866-6363 FAX 408-866-6364 EMAIL [info@listlabs.com](mailto:info@listlabs.com)  
WEBSITE [www.listlabs.com](http://www.listlabs.com)

Product #520  
Lot #5209A3  
Release Date: October 2018  
Recommended Retest Date: October 2021

CERTIFICATE OF ANALYSIS  
SNAPTide® (o-Abz/Dnp)  
Peptide Substrate for Botulinum Neurotoxin Type A  
Lot #5209A3

**Contents**

Each vial of SNAPTide® (o-Abz/Dnp) (U.S. Patent #6,504,006), a botulinum neurotoxin type A (BoNT/A) substrate, contains 200 nmoles of lyophilized peptide. This peptide is intramolecularly quenched by fluorescence resonance energy transfer (FRET). The N-terminally-linked fluorophore is o-aminobenzoic acid (o-Abz) and the acceptor chromophore is a 2,4-dinitrophenyl group (Dnp). This lyophilized powder is stoppered under vacuum. It is recommended that it be stored at -20°C, protected from light.

**Reconstitution**

A small amount of peptide has been lyophilized in each vial. During lyophilization and transportation, this material may be distributed throughout the vial. Since it is common practice to reconstitute peptide in a small volume of solvent, visually locate the powder and, if necessary, shake it to the bottom of the vial prior to adding the solvent. It is recommended that initial stock solutions be made in DMSO to ensure total recovery of the product from the vial. Cover the vial with foil to protect the peptide from light.

**Concentration**

Concentration is determined from the absorbance at 363 nm using the molar absorption coefficient of 15,900 M<sup>-1</sup>cm<sup>-1</sup> for Lys(Dnp).

**Analysis**

The peptide is >95% pure as determined by reverse phase HPLC. The expected molecular weight was obtained by mass spectrometry.

**Assay Conditions and Parameters for Utilizing SNAPTide® (o-Abz/Dnp) FRET Peptide**

**SNAPTide® (o-Abz/Dnp), Product #520**

Prepare a 5 mM stock solution of this peptide in DMSO as follows: Add 40 µl of DMSO to a vial containing 200 nmoles of peptide. Cover the vial with foil to protect from light, and store frozen at -20°C.

The FRET assays are performed using HEPES buffers prepared by titrating the free acid form of HEPES with the potassium salt form of HEPES. For assays with botulinum neurotoxin Type A (BoNT/A) holotoxin, the SNAPTide® stock solution is diluted using 20 mM HEPES, pH 8.0, prior to use. For assays with BoNT/A Light Chain, the stock solution should be diluted in the hydrolysis buffer, described in the section below. When using a 96-well plate and a final volume of 250 µl/well, a 250 µM stock solution is convenient to use. The final concentration of SNAPTide® to be used is typically between 5 µM and 10 µM/well, depending on the instrumentation and experiment. Since DMSO inhibits cleavage, final concentrations must be less than 2% of the total volume. For SNAPTide® (o-Abz/Dnp), Product #520, any concentration of ZnCl<sub>2</sub> in the BoNT/A Light Chain hydrolysis buffer inhibits cleavage.

These FRET assays are run at 37°C. Excitation wavelength is 320 nm and emission is 420 nm. There is a linear dependence of fluorescence intensity on concentration of totally cleaved substrate up to 30 µM SNAPTide® (o-Abz/Dnp).

When measuring kinetic parameters such as the K<sub>m</sub> and V<sub>max</sub> for this FRET substrate, the data must be corrected for a phenomenon known as the "inner filter effect". This effect, as well as a method to determine an appropriate correction factor, is explained in the paper by Liu *et al.* (1999) in *Analytical Biochemistry*, **267**, 331-335. The correction method uses an unquenched fluorophore for comparison. Since the fluorescence efficiency for the free o-Abz is higher than that for o-Abz when it is bound to the peptide, the use of product #529, Unquenched Calibration Peptide for SNAPTide® 520, in the place of the free o-Abz, is suggested. This peptide contains the o-Abz bound to the N-terminal cleaved fragment of SNAPTide®.

(continued)

**Botulinum Neurotoxin Type A (BoNT/A) Holotoxin, Product #130/#9130**

It is recommended to reconstitute this protein with the reduction buffer, 20 mM HEPES, pH 8.0, containing 5 mM DTT, 0.3 mM ZnSO<sub>4</sub> and 0.1% TWEEN 20. In order to activate BoNT/A holotoxin it must be reduced by incubation for 30 minutes at 37°C immediately following reconstitution in this buffer. Use reduced toxin as soon as possible. Concentrations of BoNT/A holotoxin between 2 nM and 10 nM can be used depending on the instrumentation and experiment. The TWEEN 20 in the reduction buffer is essential for recovery of the BoNT/A holotoxin from the vial. It is possible to substitute 1 mg/ml BSA for the TWEEN 20. The reaction buffer for the hydrolysis of SNAPtide® using BoNT/A holotoxin is 20 mM HEPES, pH 8.0, containing 0.75 mM ZnSO<sub>4</sub>, 1.25 mM DTT and 0.1% TWEEN 20.

**Botulinum Neurotoxin Type A Light Chain, Recombinant, Product #610A**

For the reconstitution of BoNT/A Light Chain and for the hydrolysis reaction of SNAPtide® with BoNT/A Light Chain, use the hydrolysis buffer 50 mM HEPES, pH 7.4, containing 0.05% TWEEN 20. BoNT/A Light Chain does not require reduction. Concentrations of BoNT/A Light Chain between 2 nM and 10 nM can be used depending on the instrumentation and experiment. The addition of 0.05% TWEEN 20 or 1 mg/ml BSA is beneficial to the stability and storage of reconstituted BoNT/A Light Chain at -20°C.

Note: Recent studies analyzing the effect of the osmolyte, trimethylamine N-oxide (TMAO), on the rate of cleavage of SNAPtide®, Prod #520 with BoNT/A Light Chain indicate that the rate decreases as the concentration of the TMAO increases. The opposite effect is observed with SNAPtide®, Prod #521.

**Related Products**

- Product #529:** Unquenched Calibration Peptide for SNAPtide® 520 Substrate for *C.botulinum* Type A Neurotoxin. A calibration peptide which is the cleavage product of SNAPtide® containing only the o-Abz at the N-terminal; it can be used to convert relative fluorescence units (RFU) to nmoles of cleaved substrate.
- Product #521:** SNAPtide® Peptide Substrate (FITC/DABCYL) for *C.botulinum* Type A Neurotoxin. (U.S. Patent #6,504,006)
- Product #528:** Unquenched Calibration Peptide for SNAPtide® 521 Substrate for *C.botulinum* Type A Neurotoxin. A calibration peptide which is the cleavage product of SNAPtide® containing only the FITC at the N-terminal; it can be used to convert relative fluorescence units (RFU) to nmoles of cleaved substrate.
- Product #523:** SNAPtide® Peptide Substrate fIP6 (DABCYL/5-IAF) for *C.botulinum* Type A Neurotoxin. (U.S. Patent Pending #61/252,675)
- Product #525:** Control Peptide for SNAPtide® 520 Substrate for *C.botulinum* Type A Neurotoxin. The control peptide for SNAPtide® 520 contains two substitutions and is not a substrate for BoNT/A, however, since it contains all of the sites for non-specific cleavage found in SNAPtide® 520, it is an ideal control peptide.

For further information regarding this FRET peptide and related products, click on the Posters tab on our website, <https://www.listlabs.com/blog/list-labs-scientific-posters-available-on-the-website/>.

**Handling**

This product is not known to be hazardous. Good laboratory technique should be employed in the safe handling of this product. Wear appropriate laboratory attire including a lab coat, gloves and safety glasses. Nitrile gloves are recommended when handling lyophilized material.

This product is intended for research purposes only. It is not intended for use in humans or as a diagnostic agent. List Biological Laboratories, Inc. is not liable for any damages resulting from the misuse or handling of this product.

**FOR RESEARCH PURPOSES ONLY. NOT FOR USE IN HUMANS.**

Production: KS Date: 10/30/18 Management: NS Date: 10/30/18 QA/QC: YLP Date: 10/30/18