

FITC AND BIOTIN CONJUGATES OF CHOLERA TOXIN B SUBUNIT

Conjugates of cholera toxin and cholera toxin B subunit (choleragenoid) have been employed as histochemical markers in both axoplasmic transport and neuronal connectivity studies.¹ The basis of this lies in the binding specificity of choleragenoid to GM1 ganglioside receptors on neuronal cell surfaces.² Cholera toxin B subunit retains the full binding capacity of the intact toxin, yet has the advantage of being completely nontoxic.

List Biological Laboratories, Inc. offers cholera toxin B subunit conjugated to fluorescein (CTB FITC), a sensitive fluorescent probe suitable for use in flow cytometry and microscopy,³ and biotinylated cholera toxin B subunit (CTB biotin) for high affinity cytochemistry for microscopy.^{4,5} Each of these conjugates are supplied as aseptically packaged lyophilized powders. A detailed lot analysis and instructions on reconstitution, storage and handling are provided with each shipment.

Goat anti-choleragenoid, suitable for neutralization and binding assays or immunohistochemical studies in conjunction with unconjugated cholera toxin B subunit or either of the above products, is also available from List Biological Laboratories, Inc.^{6,7,8} This product is provided as an aseptically packaged lyophilized powder and contains 0.1% sodium azide as a preservative.

The above products are intended for research purposes only and are not for use in humans or as diagnostic agents.

Ordering Information

Product No.	Description	Size
104	Cholera Toxin B Subunit (Low Salt)	0.5 mg
106	Cholera Toxin B Subunit Fluorescein Isothiocyanate Conjugate (CTB FITC)	0.2 mg
112	Cholera Toxin B Subunit Biotin Conjugate (CTB Biotin)	0.2 mg
703	Goat Anti-Choleragenoid Serum	0.1 ml

See how others have used List Labs' products on our citations page: <https://www.listlabs.com/citations>

References

1. Grafstein B, Forman D. Intracellular transport in neurons. *Physiol. Rev.* 1980; 60(4):1168-1283. [PMID:6159657](#)
2. Van Heyningen WE. [Gangliosides as membrane receptor for tetanus toxin, cholera toxin and serotonin.](#) *Nature* 1974; 249:415-417.
3. Mishima H, Sears M, Bausher L, Gregory D. Ultracytochemistry of cholera-toxin binding sites in ciliary processes. *Cell Tissue Research.* 1982; 223(2):241-253. [PMID:7066973](#)
4. Bayer EA, Skatelsky E, Wilchek M. The Avidin-Biotin Complex in Affinity Cytochemistry. *Meth. Enz.* 1979; 62(D):308-315. [PMID:440114](#)
5. Asou H, Brunngraber EG, Jeng I. Cellular localization of G_{M1}-ganglioside with biotinylated cholera toxin and avidin peroxidase in primary cultured cells from rat brain. *J. Histochem. Cytochem.* 1983; 31(12):1375-1379. [PMID:6195214](#)
6. Hayakawa T, Zheng JQ, Seki M. Direct parabrachial nuclear projections to the pharyngeal motoneurons in the rat: an anterograde and retrograde double-labeling study. *Brain Research.* 1999; 816(2):364-374. [PMID:9878830](#)
7. Gaillard F, Letang J, Frappe I, Gaillard A. Laminar distribution of isocortical neurons projecting to occipital grafts in neonate and adult rats. *Experimental Neurology.* 2000; 162(1):225-233. [PMID:10716903](#)
8. Ruigrok TJ, van Touw S, Coulon P. Caveats in Transneuronal Tracing with Unmodified Rabies Virus: An Evaluation of Aberrant Results Using a Nearly Perfect Tracing Technique. *Front Neural Circuits.* 2016; 10:46. [PMID:27462206.](#)

©1983 LBL, Inc. Rev. 09/2017